

JOURNAL OF ALGEBRA 63, 484–493 (1980)

Simple Lie Algebras with Classical Reductive Null Component

THOMAS B. GREGORY

*Department of Mathematics, Ohio State University, Mansfield, Ohio 44906**Communicated by N. Jacobson*

Received May 20, 1978; revised July 2, 1979

INTRODUCTION

In [2], Kac proposed a program to classify the simple finite-dimensional Lie algebras over an algebraically closed field of characteristic greater than three. The first (and apparently most difficult) step in this program is finding an appropriate maximal subalgebra L_0 of a given Lie algebra L : it is required that L_0 determine a filtration $L = L_{-k} \supseteq \cdots \supseteq L_0 \supseteq \cdots$ in which L_0/L_1 is the direct sum of simple classical algebras and a possibly nontrivial center. Candidates for L_0 have been proposed by Kostrikin [4] and Wilson [12] and work is progressing in that area. The next step in the program is to determine the associated graded algebra $\text{gr } L$ when L_0/L_1 is known. The final step is to find the filtered Lie algebras with which a particular graded Lie algebra is associated.

Wilson [11], Kostrikin and Šafarevič [6], and Kac [2, 3] have made significant progress in carrying out the last two steps. In connection with the next-to-last step, Kac [2, Theorem 3] proves the following theorem: Let $L = \bigoplus G_i$ be a finite-dimensional nontrivial transitive graded Lie algebra over an algebraically closed field of characteristic $p > 3$ satisfying the following conditions:

- (a) The subalgebra G_0 is isomorphic to the direct sum of its center and some simple Lie algebras of classical type.
- (b) The G_0 -module G_{-1} is a p -representation.

Then G is isomorphic either to a Lie algebra of Cartan type or to a Lie algebra of classical type with one of the standard gradations, or to a Lie algebra of type $A_{k,p-1}^z$ (see the definition following Proposition 1.5 of [2]). The present author has shown that if it is only assumed that $L = G_{-1} \oplus G_0 \oplus \cdots \oplus G_k$ is a simple Lie algebra over an algebraically closed field of characteristic $p > 3$ such that condition (a) above holds, then G is isomorphic either to a Lie algebra of Cartan type, or to a Lie algebra of classical type with one of the standard gradations, or to a Lie algebra of type $A_{k,p-1}^z$.

In [5], Kostrikin solved a similar problem: he determined the simple graded Lie algebras whose zero component is isomorphic to W_1 and whose -2 component is zero. More recently, Kuznecov [7] showed that there are no simple graded Lie algebras whose null component is solvable.

0. NOTATION

Throughout this paper, L will denote a simple, graded Lie algebra over an algebraically closed field of characteristic $p > 3$. We will assume that L has the particular form

$$L = G_{-1} \oplus G_0 \oplus \cdots \oplus G_k, \quad (1)$$

where $G_k \neq \{0\}$, and where, as usual, the G_i are subspaces such that

$$[G_i, G_j] \subseteq G_{i+j} \quad \text{whenever} \quad -1 \leq i + j \leq k, \quad (2)$$

and

$$[G_i, G_j] = \{0\}, \quad \text{otherwise.} \quad (3)$$

Since $[G_0, G_0] \subseteq G_0$, G_0 is a subalgebra. We assume that G_0 is isomorphic to the direct sum of its center and some simple Lie algebras of classical type:

$$G_0 = Z \oplus G_{0,1} \oplus G_{0,2} \oplus \cdots \oplus G_{0,b} = Z \oplus [G_0, G_0], \quad (4)$$

where Z may be zero, and $b \geq 1$. Let

$$\alpha_1, \dots, \alpha_\rho \quad (5)$$

be a system of simple roots of $[G_0, G_0]$. Let e_i , f_i , and h_i be vectors in the spaces associated with the roots α_i , $-\alpha_i$, and zero, respectively, such that

$$[e_i, f_i] = h_i, \quad [e_i, h_i] = 2e_i, \quad \text{and} \quad [f_i, h_i] = -2f_i; \quad (6)$$

also,

$$[e_{i_1}, f_{i_2}] = 0 \quad (7)$$

if i_1 differs from i_2 . In view of (2), we have that $[G_0, G_j] \subseteq G_j$ for each j . The Jacobi identity in L shows that for $g_0, g'_0 \in G_0$ and $g_j \in G_j$, we have

$$[g_j, [g_0, g'_0]] = [[g_j, g_0], g'_0] - [[g_j, g'_0], g_0];$$

hence, if we take ad_{G_j} to mean the restriction to G_j of the adjoint representation

of L in itself, then $g_0 \rightarrow \text{ad}_{G_j} g_0$ gives a representation of G_0 in G_j . When we speak of a weight vector of G_j , we will mean a weight vector with respect to this representation of G_0 in G_j . (When $j = 0$, we consider the terms "weight vector" and "root vector" to be synonymous.) For each t , $1 \leq t \leq b$, we fix a highest-root vector r_t of $G_{0,t}$; i.e., r_t is a root vector of $G_{0,t}$ which is annihilated by all e_i .

The particular nonzero scalar multiple of a vector will often be of little interest in an argument. We have therefore found it convenient to use the symbol " \doteq ," which will mean "is a nonzero scalar multiple of."

1. THE MAIN RESULT

Our aim is to prove the following

THEOREM. *Let $L = G_{-1} \oplus G_0 \oplus \cdots \oplus G_k$ be a simple, finite-dimensional, graded Lie algebra over an algebraically closed field of characteristic $p > 3$, and suppose that G_0 is isomorphic to the direct sum of its (possibly trivial) center and some simple Lie algebras of classical type. Then G is isomorphic either to a Lie algebra of Cartan type or to a Lie algebra of classical type with one of the standard gradations, or to a Lie algebra of type A_{kp-1}^z .*

Remark. This theorem generalizes the theorem in Chapter II of [6], where the dimension of G_{-1} is assumed to be less than $p - 1$ and the representation of G_0 in G_{-1} induced by the bracket operation of L is assumed to be an irreducible p -representation; also, as noted in the Introduction, this theorem generalizes Theorem 3 of [2] (for L as in (1)).

2. SIMPLE PROPERTIES OF SIMPLE LIE ALGEBRAS WITH GRADATION

We begin by showing that G_{-1} and G_k are nontrivial, irreducible G_0 -modules, and that $[G_{-1}, x]$ is nonzero for any (nonzero) element x of L which is not contained in G_{-1} . Graded Lie algebras satisfying this latter condition are often called "transitive," while those in which G_{-1} is an irreducible G_0 -module are usually referred to as "irreducible" (see, for example, Section 1 of [2].)

LEMMA 1. *If i is greater than -1 , then no nonzero element of G_i is annihilated by G_{-1} . Furthermore, G_{-1} is a nontrivial, irreducible G_0 -module.*

Proof. First of all, we note that if x is a nonzero element of L which is annihilated by G_{-1} , then

$$I = \langle x \rangle + \sum_{j_1, \dots, j_n \geq 0} [\cdots [x, G_{j_1}], \dots, G_{j_n}] \quad (8)$$

is a nonzero ideal of L . It suffices to show that I is stable under $\text{ad } G_{-1}$; one easily shows $[[\cdots [x, G_{j_1}], \dots, G_{j_n}], G_{-1}] \subseteq I$ by the Jacobi identity and induction on n . If $x \in G_i \setminus \{0\}$ for some $i > -1$ and $[G_{-1}, x] = \{0\}$, then $I \subseteq G_i \oplus \cdots \oplus G_k \neq L$, so that I would be a proper ideal of the simple algebra L . This contradiction proves the first assertion of the lemma.

Now let M be a nonzero G_0 -submodule of G_{-1} , and let x be a nonzero element of M . By (3), $[G_{-1}, x] = \{0\}$. Let I be defined as in (8). Then $I \neq \{0\}$, so $I = L$, since L is simple. Because

$$I \cap G_{-1} = \langle x \rangle + \sum_{n \geq 0} [\cdots [x, \underbrace{G_0, \dots, G_0}_{n \text{ copies}}], G_{-1}]$$

it is clear that $M \supseteq I \cap G_{-1}$. Therefore, we have

$$G_{-1} \supseteq M \supseteq I \cap G_{-1} = L \cap G_{-1} = G_{-1},$$

so that $M = G_{-1}$. Since M was any nonzero G_0 -submodule of G_{-1} , it follows that G_{-1} is an irreducible G_0 -module.

If G_{-1} were a trivial G_0 -module (i.e., if $[G_{-1}, G_0] = \{0\}$), then $[L, L] = \sum_{-1 \leq i, j \leq k} [G_i, G_j] \subseteq G_0 \oplus \cdots \oplus G_k \neq L$, so that $G_0 \oplus \cdots \oplus G_k$ would be a proper ideal of L . This contradiction shows that G_{-1} must be a nontrivial G_0 -module, and completes the proof of the lemma.

LEMMA 2. G_k is an irreducible G_0 -module. Furthermore, for $0 \leq i \leq k$, we have that $[G_{-1}, G_i] = G_{i-1}$.

Proof. For any nonzero x in G_k ,

$$J = \langle x \rangle + \sum_{\substack{i \geq 0 \\ j \geq 0 \\ i+j \geq 1}} [\cdots [\underbrace{[x, G_0, \dots, G_0]}_{i \text{ copies}}, \underbrace{G_{-1}, \dots, G_{-1}}_{j \text{ copies}}]]$$

is a nonzero ideal of L , since by the Jacobi identity and induction on j , each summand in the sum on the right above is stable under G_n , $n \geq 1$. Since L is simple, $J = L$. Because

$$J \cap G_k = \langle x \rangle + \sum_{i \geq 0} [\cdots [x, \underbrace{G_0, \dots, G_0}_{i \text{ copies}}], G_k]$$

it follows, as in the proof of Lemma 1, that for M a nonzero G_0 -submodule of G_k and x a nonzero element of M , we have for J defined as above in terms of this x that

$$G_k \supseteq M \supseteq J \cap G_k = L \cap G_k = G_k.$$

Therefore, $M = G_k$, and G_k is seen to be an irreducible G_0 -module.

Finally, for any j such that $0 \leq j \leq k+1$, we have

$$G_{k-j} = G_{k-j} \cap L = G_{k-j} \cap J = \sum_{i \geq 0} [\cdots [\underbrace{[x, G_0], \dots, G_0]_{i \text{ copies}}, \underbrace{G_{-1}, \dots, G_{-1}]_{j \text{ copies}}}]$$

In particular,

$$G_{k-j} = [G_{k-(j-1)}, G_{-1}],$$

which is just the second assertion of the lemma.

LEMMA 3. *If for some m and n such that $0 \leq m, n \leq k$ we have $[G_{-1}, G_m] = G_{m-1}$, $[G_m, G_n] = \{0\}$, and $[G_{-1}, G_n] = G_{n-1}$, then $[G_{m-1}, G_n] = [G_m, G_{n-1}]$.*

Proof. In view of the Jacobi identity, we have

$$[[G_m, G_{-1}], G_n] \subseteq [[G_m, G_n], G_{-1}] + [G_m, [G_{-1}, G_n]].$$

By our assumptions, the subspace on the left is equal to $[G_{m-1}, G_n]$, the first summand on the right is equal to zero, and the second summand on the right is equal to $[G_m, G_{n-1}]$; hence, $[G_{m-1}, G_n] \subseteq [G_m, G_{n-1}]$. Inclusion in the other direction holds by symmetry, and the lemma follows.

LEMMA 4. G_k is a nontrivial G_0 -module.

Proof. By (3), we have

$$[G_i, G_j] = \{0\} \quad \text{for } i+j = k+1,$$

so, in view of the second assertion of Lemma 2, we have by repeated application of Lemma 3 that

$$[G_0, G_k] = [G_1, G_{k-1}] = \cdots = [G_i, G_{k-i}] = \cdots = [G_k, G_0].$$

Hence, if G_k were a trivial G_0 -module, then all of the above subspaces would be zero. In view of (1), we would then have

$$[L, L] = \sum_{-1 \leq i, j \leq k} [G_i, G_j] \subseteq G_{-1} \oplus \cdots \oplus G_{k-1},$$

and the sum on the right above would be a proper ideal of L , contradicting our assumption that L is simple. Q.E.D.

LEMMA 5. *For each j such that $-1 \leq j \leq k$, we have $G_j \neq \{0\}$.*

Proof. We assumed at the outset (1) that $G_k \neq \{0\}$. If $G_j = \{0\}$ for some j such that $-1 \leq j < k$, then $\bigoplus_{n=j+1}^k G_n$ would be an ideal of L , contradicting the simplicity of L .

3. HIGHEST-WEIGHT VECTORS IN G_{-1}

If the representation of $[G_0, G_0]$ in G_{-1} is a p -representation, then one can conclude that G_{-1} contains a highest-weight vector for this representation. This section is devoted to showing that even when we do not assume that the representation is restricted, a highest-weight vector must nonetheless exist.

LEMMA 6. *For each j such that $-1 \leq j \leq k$, the subspace G_j contains a nonzero weight vector for the representation of G_0 in G_j .*

Proof. Since we have assumed that G_j is a finite-dimensional vector space over an algebraically closed field, this lemma follows from Lemma 5 and the commutativity of the Cartan subalgebra of G_0 .

LEMMA 7. *For $-1 \leq i \leq k-1$, G_i is spanned by weight vectors of the form $[u, w]$, where u is a weight vector of G_{-1} , and w is a weight vector of G_{i+1} . G_k is spanned by weight vectors, also.*

Proof. By Lemma 6, G_{-1} contains a nonzero weight vector x for the representation of G_0 in G_{-1} . According to Lemma 1, G_{-1} is irreducible as a G_0 -module, so G_{-1} must coincide with the G_0 -submodule S of G_{-1} which is generated by x . Since G_0 is spanned by root vectors and the bracket of a root vector with a weight vector is a weight vector (or zero), it follows that G_{-1} ($= S$) is spanned by weight vectors. Similarly, by Lemmas 6 and 2, G_k is spanned by weight vectors. Since the bracket of two weight vectors is again a weight vector (or zero), the first assertion of Lemma 7 now follows from the second assertion of Lemma 2 and induction. Q.E.D.

LEMMA 8. *G_{-1} contains a weight vector v^- which is annihilated by all e_i .*

Proof. Fix t . Then $r_t = \sum_{j=1}^m [u_j, w_j]$ by Lemma 7, where u_j is a weight vector of G_{-1} and w_j is a weight vector of G_1 . Since the root-space decomposition of $G_{0,t}$ is direct (see, for example, Sections I.5 and I.6 of [8]), and since the root space corresponding to the highest root is one dimensional (Lemma II.3.2 of [8]), we can assume that for every j , $[u_j, w_j]$ is either zero or a nonzero scalar multiple of r_t . Let j_0 be an integer such that $[u_{j_0}, w_{j_0}] \neq 0$. Then we set $v = u_{j_0}$ and $c = w_{j_0}$, and we have

$$[v, c] = r_t.$$

By Lemma 1, $[G_{-1}, r_t] \neq \{0\}$; hence, there exists a weight vector $y \in G_{-1}$ such that $[r_t, y] \neq 0$. Suppose there were an infinite sequence of simple-root vectors e_{s_1}, e_{s_2}, \dots of G_0 such that

$$[\cdots [[r_t, y], e_{s_1}], \dots, e_{s_n}] = [r_t, [\cdots [y, e_{s_1}], \dots, e_{s_n}]] \quad (9)$$

were unequal to zero for all n . Since $r_n \doteq [v, c]$ and G_{-1} is Abelian, this would imply that

$$[[v, c], [\cdots [y, e_{s_1}], \dots, e_{s_n}]] = [v, [c, [\cdots [y, e_{s_1}], \dots, e_{s_n}]]]$$

were unequal to zero, so that

$$[c, [\cdots [y, e_{s_1}], \dots, e_{s_n}]] \neq 0$$

for all n . But these are root vectors of G_0 , so we would have an infinite sequence of roots of G_0 , each obtained from the previous root by the addition of a simple (positive) root. This situation is known to be impossible when G_0 is classical (for the proof for prime characteristic, see [8, p. 31]). Hence, no such infinite sequence of nonzero vectors as (9) can exist; i.e., any sequence (9) must terminate in a vector which is annihilated by every simple-root vector e_i . We take v^+ to be such a terminal vector. Q.E.D.

4. PROOF OF THE MAIN RESULT

The present author's original proof of the main result (for $k > 1$ and $G_0 = G_{0,1}$) proceeded along the following lines: if the difference in the weights of the highest- and lowest-weight vectors of G_{-1} were greater than the highest root of G_0 , then the difference between the highest and lowest roots of G_0 would be greater than twice the highest root; i.e., we arrive at a contradiction. Using the fact that

$$v^+ \doteq [v^-, r_1], \quad (10)$$

where v^- is a lowest-weight vector of G_{-1} , one has that $\text{ad}_{G_{-1}} r_1$ is a rank one transformation of G_{-1} and $(\text{ad}_{G_0} r_1)^2 \neq 0$, so that Theorem 1 of [9] implies the result. Also, one can use (10) to show that G_0 must be A_n or C_n and the representation Γ of G_0 on G_{-1} must be the natural representation of G_0 on G_{-1} . Then Theorem 2 of Chapter III of [6] implies the result. Finally, (10) can be used to show that Γ is restricted, and one can then use Theorem 3 of [2] to get the result. The present author also proved the main result for $k = 1$ and G_0 simple; the referee pointed out that this case had already been dealt with in [2, 9].

The author wishes to thank the referee for suggesting the shorter reduction to Theorem 3 of [2] below. Also, thanks are due to Professor George B. Seligman for his help in generalizing the result and his formulation of the referee's ideas.

Proof of Theorem (Referee; Seligman). Since the choice of a root system (5) for $[G_0, G_0]$ was arbitrary, Lemma 8 shows that for every classical Cartan decomposition of $[G_0, G_0]$ with associated ordering of the roots, there is a nonzero weight vector v^+ in G_{-1} that is annihilated by all root vectors belonging to positive roots.

If e_i is a positive-root vector, then $(\text{ad}_{G_0} e_i)^p = 0$. It follows that e_i^p is in the center of the universal enveloping algebra $U(G_0)$, so acts as a scalar on G_{-1} by Schur's lemma. But e_i annihilates v^+ , so e_i^p does, too; the scalar must therefore be zero, and e_i^p annihilates G_{-1} .

Now one can use the Weisfeiler-Kac linear function $l(x) = (\chi_{G_{-1}}(x^p - x^{[p]}))^{1/p}$ (see [10]): this annihilates all root vectors by the above remarks, based on Lemma 8, and $[G_0, G_0]$ is spanned by root vectors relative to classical Cartan decompositions. (For this, fix a classical Cartan decomposition. Two applications of the above give (i) $l(x) = 0$ if x is a positive-root vector; (ii) $l(x) = 0$ if x is a negative-root vector. It remains only to show that l vanishes on the Cartan subalgebra; but each of the automorphisms $\exp(\text{ad}_{G_0} e_i)$ maps classical Cartan decompositions to classical Cartan decompositions. Thus,

$$l(f_i^{\exp(\text{ad}_{G_0} e_i)}) = 0;$$

but

$$f_i^{\exp(\text{ad}_{G_0} e_i)} = f_i + h_i + k_i e_i,$$

where k_i is an element of the base field. It follows from (i), (ii), and the above that $l(h_i) = 0$.) Thus, $x^p - x^{[p]}$ acts as the scalar 0 for each $x \in [G_0, G_0]$ so G_{-1} is a p -module of $[G_0, G_0]$.

Now let y be any nonzero element of Z . Then by Schur's lemma, $\text{ad}_{G_{-1}} y = \lambda I$, where I is the identity transformation of G_{-1} and λ is an element of the base field. By the first assertion of Lemma 1, $\lambda \neq 0$. Similarly, if y' is another element of Z , then $\text{ad}_{G_{-1}} y' = \mu I$, where μ is an element of the base field. Then $\text{ad}_{G_{-1}}(y' - (\mu/\lambda)y) = 0$, so, again by Lemma 1, $y' - (\mu/\lambda)y = 0$; i.e., $Z = \langle y \rangle$.

Let $z = -(1/\lambda)y$. Then $Z = \langle z \rangle$, and $\text{ad}_{G_{-1}} z = -I$. We make Z into a p -algebra by defining $z^{[p]} = z$. Then

$$(\text{ad}_{G_{-1}} z)^p = (-I)^p = -I = \text{ad}_{G_{-1}} z = \text{ad}_{G_{-1}} z^{[p]},$$

so G_{-1} is a p -module of Z , also.

The restriction of the representation Γ of G_0 in G_{-1} to each of the direct summands (4) of G_0 is a restricted representation. It follows that Γ is restricted. Our main result now follows from Theorem 3 of [2].

APPENDIX: NOTATION

L	a simple, graded Lie algebra over an algebraically closed field of characteristic $p > 3$
G_i	i th subspace in the gradation of L
z	vector spanning the center of G_0
$G_{0,t}$	t th simple Lie algebra summand of G_0
k	highest index of a gradation subspace of L
b	number of simple Lie algebra summands of G_0
α_i	i th simple root in an arbitrary ordering of simple roots of G_0
e_i	vector spanning the space associated with the root α_i
f_i	vector spanning the space associated with the root $-\alpha_i$
h_i	vector equal to the bracket of e_i with f_i
g_i	arbitrary vector in G_i
g'_i	arbitrary vector in G_i
\doteq	symbol meaning "is a nonzero scalar multiple of"
r_t	root vector of $G_{0,t}$ which is annihilated by e_i , $i = 1, \dots, \rho$
v^+	weight vector of G_{-1} which is annihilated by e_i , $i = 1, \dots, \rho$
$\chi_{G_{-1}}$	homomorphism from the center of the universal enveloping algebra of G_0 to the base field defined in [10]
Γ	representation of G_0 in G_{-1} induced by the bracket operation in L
ρ	number of simple roots of G_0

ACKNOWLEDGMENTS

The present article is a generalization of the author's dissertation [1], written under the gracious and ever helpful guidance of Professor George B. Seligman, to whom the author expresses his deepest appreciation. Both Professor Seligman and Professor Robert Lee Wilson provided many valuable suggestions which improved that work considerably, and Professor Seligman has continued his most generous assistance in the preparation of this article.

REFERENCES

1. T. B. GREGORY, Simple Lie algebras with classical simple null component, Dissertation, Yale University, 1977.
2. V. G. KAC, The classification of the simple Lie algebras over a field with non-zero characteristic, *Izv. Akad. Nauk SSSR Ser. Mat.* **34** (1970), 385-408; Engl. transl., *Math. USSR Izv.* **4** (1970), 391-413.

3. V. G. KAC, Description of filtered Lie algebras with which graded Lie algebras of Cartan type are associated, *Izv. Akad. Nauk SSSR, Ser. Mat.* **38** (1974), 800–838; Engl. transl., *Math USSR Izv.* **8** (1974), 801–835.
4. A. I. KOSTRIKIN, Squares of adjoint endomorphisms in simple Lie p -algebras, *Izv. Akad. Nauk SSSR, Ser. Mat.* **31** (1967), 445–487; Engl. transl., *Math USSR Izv.* **1** (1967), 434–473.
5. A. I. KOSTRIKIN, Irreducible graded Lie algebras with the component $L_0 = W_1$ (Russian), *Ural. Gos. Univ. Mat. Zap.* **7** (1969/1970), tetrad' 3, 92–103.
6. A. I. KOSTRIKIN AND I. R. ŠAFAREVIČ, Graded Lie algebras of finite characteristic, *Izv. Akad. Nauk SSSR, Ser. Mat.* **33** (1969), 251–322; Engl. transl. *Math USSR Izv.* **3** (1969), 237–304.
7. M. K. KUZNECOV, Simple modular Lie algebras with a solvable maximal subalgebra (Russian), *Mat. Sb.* **101** (143), 1976, 77–86.
8. G. B. SELIGMAN, "Modular Lie Algebras," *Ergebnisse der Math.*, Band 40, Springer-Verlag, New York/Berlin, 1967.
9. B. JU. WEISFEILER AND V. G. KAC, Exponentials in Lie algebras of characteristic p , *Izv. Akad. Nauk SSSR, Ser. Mat.* **35** (1971), 762–788.
10. B. JU. WEISFEILER AND V. G. KAC, Irreducible representations of Lie p -algebras, *Funktsional Anal. i Prilozhen.* **5**, No. 2 (April–June, 1971), 28–36.
11. R. L. WILSON, A structural characterization of the simple Lie algebras of generalized Cartan type over fields of prime characteristic, *J. Algebra* **40** (1976), 418–465.
12. R. L. WILSON, Recent progress in the classification of simple Lie algebras over algebraically closed fields of prime characteristic, mimeograph, Rutgers University, 1975.